

Yale University

EliScholar – A Digital Platform for Scholarly Publishing at Yale

Cowles Foundation Discussion Papers

Cowles Foundation

6-1-1995

Unit Root Tests

Peter C.B. Phillips

Follow this and additional works at: <https://elischolar.library.yale.edu/cowles-discussion-paper-series>



Part of the [Economics Commons](#)

Recommended Citation

Phillips, Peter C.B., "Unit Root Tests" (1995). *Cowles Foundation Discussion Papers*. 1347.
<https://elischolar.library.yale.edu/cowles-discussion-paper-series/1347>

This Discussion Paper is brought to you for free and open access by the Cowles Foundation at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Cowles Foundation Discussion Papers by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

COWLES FOUNDATION FOR RESEARCH IN ECONOMICS
AT YALE UNIVERSITY

Box 2125, Yale Station
New Haven, Connecticut 06520

COWLES FOUNDATION DISCUSSION PAPER NO. 1104

Note: Cowles Foundation Discussion Papers are preliminary materials circulated to stimulate discussion and critical comment. Requests for single copies of a Paper will be filled by the Cowles Foundation within the limits of the supply. References in publications to Discussion Papers (other than mere acknowledgment by a writer that he has access to such unpublished material) should be cleared with the author to protect the tentative character of these papers.

UNIT ROOT TESTS

Peter C. B. Phillips

June 1995

Unit Root Tests

by

Peter C. B. Phillips*

*Cowles Foundation for Research in Economics
Yale University*

Abstract

Classical and Bayesian unit root test procedures are reviewed, with an emphasis on testing principles and recent developments. A numerical illustration and annotated references and bibliography are provided.

Keywords: Autoregressive unit root; Brownian motion; Functional central limit theorem; integrated process; LM principle; model selection; moving average unit root; nonstationarity; quasi-differencing; stationarity; stochastic trend.

JEL Classification: C22.

* Prepared as an entry for inclusion in the Encyclopedia of Statistical Sciences. The paper was keyboarded by the author in MSWord 6. Thanks go Bruce Hansen and Peter Schmidt for comments on a preliminary draft, and to the NSF for research support under grant no. SES 9122142.

ESS Update

Peter C. B. Phillips

Unit Root Tests

Autoregressive unit root; Brownian motion; difference stationarity; integrated process; model selection; nonstationarity; stochastic trend; trend stationarity.

Many observed time series display nonstationary characteristics. Some grow in a secular way over long periods of time, others appear to wander around in a random way as if they have no fixed population mean. These characteristics are especially evident in time series that represent aggregate economic behaviour (like gross domestic product), financial time series (such as indexes of stock prices), and political opinion poll series (such as presidential popularity data). Any attempt to explain or forecast series of this type requires that a mechanism be introduced to capture the nonstationary elements in the series, or that the series be transformed in some way to achieve stationarity. The problem is particularly delicate in the multivariate case, where several time series may have nonstationary characteristics and the interrelationships of these variables are the main object of study. Figure 1 graphs the monthly leading economic indicators time series for the U.S. economy over the period 1948:1 - 1994:1. Also shown in the figure is the regression line of a linear trend. The time series shows evidence of growth over time as well as a tendency to wander randomly away from the linear trend line. A successful statistical model of the time series needs to deal with both these features of the data.

One way of modeling nonstationarity* is to use deterministic trending functions like time polynomials to represent secular characteristics like growth over time. In this approach, a time series y_t is broken down into two components, one to capture trend and another to capture stationary fluctuations. A general model of this form is

$$(1) \quad y_t = h_t + y_t^s, \quad h_t = \gamma' x_t, \quad (t = 1, \dots, n),$$

where y_t^s is a *stationary time series**, x_t is an m -vector of deterministic trends* and γ is a vector of m parameters. In this case, y_t is known as a *trend-stationary* time series. The simplest example is a linear trend. Then $\gamma' x_t = \gamma_0 + \gamma_1 t$, and the time series y_t is stationary about this

deterministic linear trend. A more general example where the trends are piecewise higher order polynomials is given in equation (10) below.

Figure 1 about here

An unsatisfactory feature of trend stationary models (like the linear trend line in Figure 1) is that no random elements appear in the trending mechanism and only the stationary component is subject to stochastic shocks. Models with autoregressive *unit roots* are a simple attempt to deal with this shortcoming. In such models the trend is permitted to have both deterministic and stochastic elements. For example, in (1) the deterministic trend h_t can be retained, and the process y_t^s can be modeled as the nonstationary autoregression

$$(2) \quad y_t^s = \alpha y_{t-1}^s + u_t, \quad (t = 1, \dots, n), \quad \text{with } \alpha = 1.$$

In this model there is an autoregressive root of unity (corresponding to the solution of the characteristic equation $1 - \alpha L = 0$), and the shock u_t is stationary. *Unit root tests* usually seek to determine whether data support this model or a trend stationary alternative. In a unit root test the null hypothesis is that the autoregressive parameter $\alpha = 1$ in (2). The process y_t^s is then *difference-stationary* in the sense that the first differences $\Delta y_t^s = u_t$ are stationary. Unit root tests are typically one-sided* tests against the alternative hypothesis that $|\alpha| < 1$. Under the alternative hypothesis, the process y_t^s is stationary, and then y_t in (1) is trend stationary. Unit root tests can therefore be interpreted as tests of difference-stationarity versus trend-stationarity.

If the initial condition in (2) is set at $t = 0$, the output of the model can be written in terms of accumulated shocks as $y_t^s = \sum_{j=1}^t u_j + y_0^s$. In view of this representation, y_t^s is often called an *integrated process* of order one (written as $I(1)$). The terminology *stochastic trend* is also in common use, and is explained by the fact that y_t^s is of stochastic order $O_p(t^{1/2})$ under very general conditions, *i.e.* the variance of y_t^s is of order $O(t)$ and the standardised quantity $t^{-1/2} y_t^s$ satisfies a central limit theorem* as $t \rightarrow \infty$. The simplest example of a stochastic trend is a random

walk*. In this case, the shocks u_t are independently and identically distributed (iid) with zero mean and constant variance σ^2 . A more general case occurs when the stationary shocks u_t in (2) are generated by the linear process $u_t = C(L)\varepsilon_t$, whose innovations ε_t are iid(0, σ^2), and where $C(L)$ is a polynomial in the lag operator L for which $Ly_t = y_{t-1}$. More specifically, if

$$(3) \quad C(L) = \sum_{j=0}^{\infty} c_j L^j, \quad \sum_{j=0}^{\infty} c_j^2 < \infty, \quad \text{and} \quad C(1) \neq 0,$$

then the process u_t is covariance stationary* and has positive spectral density* at the origin, given by the expression $(\sigma^2 / 2\pi)C(1)^2$. The latter property ensures that the unit root in y_t^s does not cancel (as it would if the process u_t had a *moving average unit root*, in which case the spectral density would be zero at the origin). If the summability condition in (3) is strengthened to $\sum_{j=0}^{\infty} j^{1/2} |c_j| < \infty$, then y_t^s satisfies an invariance principle* or functional central limit theorem* (see Phillips & Solo [31] for a demonstration), and this is an important element in the development of the asymptotic theory of all unit root tests. Thus, $n^{-1/2} y_{[nr]}^s \Rightarrow B(r)$, a Brownian motion with variance $\omega^2 = \sigma^2 C(1)^2$, where $[nr]$ signifies the integer part of nr , " \Rightarrow " signifies *weak convergence**, and $r \in [0,1]$ is some fraction of the sample data. The parameter ω^2 is called the *long-run variance* of u_t .

The literature on unit root tests is vast. Most of the research has appeared since 1980, but an important early contribution came from White [39], who first recognised the vital role played by invariance principles in the asymptotic theory of time series with a unit root. The first explicit research on unit root tests dealt with Gaussian random walks and was done by Dickey & Fuller [4,5]. Solo [37], Phillips [24] and Chan & Wei [3] developed more general limit theories using invariance principles. Subsequently, an immense variety of tests have been developed, inspired in large part by the need to allow for more general processes than random walks in empirical applications. This entry covers the main principles of testing, the commonly used tests in practical work and the most recent developments.

Under certain conditions, (1) and (2) can be combined to give the regression model

$$(4) \quad y_t = \beta' x_t + \alpha y_{t-1} + u_t,$$

where β is an m -vector of deterministic trend coefficients. This formulation usually involves raising the degree of the deterministic trends to ensure that the maximum trend degrees in (4) and (1) are the same, which results in some inefficiency in the regression because there are surplus trend variables in (4). There is an alternative approach that avoid this problem of redundant variables and it will be discussed below. Asymptotic theory assumes that there exists a matrix D_n and a piecewise continuous function $X(r)$ such that $D_n^{-1} x_{[nr]} \rightarrow X(r)$ as $n \rightarrow \infty$ uniformly in $r \in [0,1]$. $X(r)$ is then the limiting trend function.

The stationary process u_t in (4) may be treated in a parametric or a non-parametric way, leading to two classes of unit root test. One relies on casting the stationary part of the process in terms of a parametric model (commonly an autoregression). The other is parametric only in its treatment of the regression coefficient α , being non-parametric with regard to the general stationary part of the process. The approach is therefore said to be semi-parametric.

The Dickey-Fuller tests and semi-parametric extensions

Let $\hat{\alpha}$ be the OLS estimator of α in (4). The Dickey-Fuller [4,5] unit root tests are based on the coefficient estimator $\hat{\alpha}$ and its regression t-ratio $t_{\hat{\alpha}}$. The basic idea of the tests is to assess whether the observed $\hat{\alpha}$ is close enough to unity to support the hypothesis of the presence of a unit root in the true data generating mechanism. Classical test procedures require a distribution theory to deliver critical values for the test statistics $\hat{\alpha}$ and $t_{\hat{\alpha}}$ under the null hypothesis that $\alpha = 1$. The finite sample distributions of these test statistics are complex and depend on unknown nuisance parameters associated with the stationary process u_t . It is therefore customary to rely on asymptotic theory, where the results are simpler and the parameter dependencies are clearly understood.

The large sample theory for $\hat{\alpha}$ and $t_{\hat{\alpha}}$ is most simply obtained using invariance principles and involves functionals of Brownian motion. In the special case where there is no deterministic

component in (4) and the shocks u_t are iid($0, \sigma^2$), the limit theory for the test statistics is as follows: $n(\hat{\alpha} - 1) \Rightarrow \left[\int_0^1 W dW \right] \left[\int_0^1 W^2 \right]^{-1}$, and $t_{\hat{\alpha}} \Rightarrow \left[\int_0^1 W dW \right] \left[\int_0^1 W^2 \right]^{-1/2}$, where W is standard Brownian motion. These limit distributions are commonly known as the Dickey-Fuller distributions, although the Brownian motion forms were not used in [4,5] and were given later in [3,24,37].

Figure 2 about here

The limit distribution of $\hat{\alpha}$ is asymmetric and has a long left tail, as shown in Figure 2. It was computed directly in [9]. In the general case where u_t is stationary, the limit has an additional bias term that depends on the autocovariance* in u_t through the nuisance parameter $\lambda = \sum_{j=1}^{\infty} E(u_0 u_j)$. This parameter and the related nuisance parameter ω^2 may be consistently estimated by *kernel** techniques, using residuals from an OLS regression on (4). If $\hat{\omega}^2$ and $\hat{\lambda}$ are such estimates, then the following statistics provide general semi-parametric tests of the unit root hypothesis (Phillips [24]), which correct for possible autocorrelation in u_t :

$$(5) \quad Z_{\alpha} = n(\hat{\alpha} - 1) - \hat{\lambda} \left(n^{-2} \sum_{t=2}^n y_{X,t-1}^2 \right)^{-1} \Rightarrow \left[\int_0^1 W_X dW \right] \left[\int_0^1 W_X^2 \right]^{-1},$$

$$(6) \quad Z_t = \hat{\sigma}_u \hat{\omega}^{-1} t_{\hat{\alpha}} - \hat{\lambda} \left\{ \hat{\omega} \left(n^{-2} \sum_{t=2}^n y_{X,t-1}^2 \right)^{1/2} \right\}^{-1} \Rightarrow \left[\int_0^1 W_X dW \right] \left[\int_0^1 W_X^2 \right]^{-1/2}.$$

In these formulae, $y_{X,t}$ is the residual from a regression of y_t on x_t , $\hat{\sigma}_u^2$ is the OLS estimator of $\sigma_u^2 = \text{var}(u_t)$, and W_X is the $L_2[0,1]$ Hilbert space projection of W onto the space orthogonal to X , viz. $W_X(r) = W(r) - \left[\int_0^1 W X \right] \left[\int_0^1 X X \right]^{-1} X(r)$.

The limit variates that appear on the right side of (5) and (6) are free from the nuisance parameters β , ω^2 , and λ , and are used to construct critical values for the tests. This is typically done by large scale simulations, since the limit distributions are non-standard. Figure 2 shows how these distributions change by stretching out the left tail as we move from a regression with no trend to a regression with a linear trend. Computerized tabulations of the critical values are given

in Ouliaris & Phillips [21] for the case of polynomial trends. In the case of the Z_α test, for instance, we reject the null hypothesis of a unit root at the 5% level if $Z_\alpha < cv(Z_\alpha; 5\%)$, the 5% critical value of the test. Both the Z_α and Z_t tests are one-sided. They measure the support in the data for a unit root against the alternative that the data are stationary about the deterministic trend x_t . When no deterministic trend appears in the model, the alternative hypothesis is just stationarity. In this case, the limit variates involve only the standard Brownian motion W . The Z_α and Z_t tests were developed in Phillips [24] and extend the original unit root tests of Dickey & Fuller based on the statistics $n(\hat{\alpha} - 1)$ and t_α . Extensions of these semi-parametric tests were obtained in [20, 22, 23, 28], and are covered by the above formulae.

To illustrate, model (4) was estimated with a linear deterministic trend for the data shown in Figure 1. The calculated values of the coefficient-based test statistics are as follows: $n(\hat{\alpha} - 1) = -7.38$; $Z_\alpha = -13.25$. The asymptotic 5% critical value of the limit distribution of the Z_α statistic is -21.21 (*c.f.* the density given by the broken line in Figure 2). These tests do not reject the null of a unit root in the time series, while allowing for the presence of a linear trend. The t-ratio test statistics are: $t_\alpha = -1.92$; $Z_t = -2.56$. The asymptotic 5% critical value of the Z_t statistic is -3.43. Again, the tests do not reject the null hypothesis of a unit root in the series. Note that the calculated values of the Dickey-Fuller statistics $n(\hat{\alpha} - 1)$ and t_α are further from the critical values than the semi-parametric statistics Z_α and Z_t . The semi-parametric corrections in the Z-tests for autocorrelation in the residual process u_t are non negligible, but in this case they do not make a difference in the outcome of the unit root tests.

The von-Neumann ratio/LM test

The von-Neumann (VN) ratio is the ratio of the sample variances of the differences and the levels of a time series. For Gaussian data this ratio leads to well known tests of *serial correlation** that have good finite sample properties. Sargan & Bhargava [34] suggested the use of this statistic for testing the Gaussian random walk hypothesis. Using non-parametric estimates

of the nuisance parameter ω^2 , it is a simple matter to rescale the VN ratio to give a unit root test for the model (1) and (2). Using a different approach and working with polynomial trends, Schmidt & Phillips [35] showed that for a Gaussian likelihood the Lagrange multiplier (LM) principle leads to a VN test, and can be generalized by using a nonparametric estimate of ω^2 .

If y_t^s were observable, the VN ratio would take the form $VN = \sum_{t=2}^n (\Delta y_t^s)^2 / \sum_{t=1}^n (y_t^s)^2$. The process y_t^s is, in fact, unobserved but may be estimated from (1). Note that, under the null hypothesis and after differences are taken, this equation is trend stationary, so that by the Grenander-Rosenblatt theorem [10, ch.7] the trend function can be efficiently estimated by an OLS regression. Let $\Delta \hat{y}_t^s = \Delta y_t - \Delta \hat{h}_t$ be the residuals from this detrending regression and let $\hat{y}_t^s = \sum_{s=2}^t \Delta \hat{y}_s^t$ be the associated estimate of y_t^s . Rescaling the von Neumann ratio then leads to the following test statistic

$$(7) \quad R_{VN} = \frac{\hat{\omega}^2}{\hat{\sigma}_u^2} \frac{n^{-1} \sum_{t=2}^n (\Delta \hat{y}_t^s)^2}{n^{-2} \sum_{t=1}^n (\hat{y}_t^s)^2} \Rightarrow \left[\int_0^1 \hat{V}_X^2 \right]^{-1}.$$

The limit process $\hat{V}_X(r)$ in (11) is a detrended *generalized Brownian bridge**, whose precise form depends on the trend h_t . In the case of a linear trend, $\hat{V}_X(r) = V(r) - \int_0^1 V$ is a demeaned version of the standard *Brownian bridge** $V(r) = W(r) - rW(1)$.

Critical values of the limit variate shown in (7) are obtained by simulation. The statistic is positive almost surely and the test is a one-sided test. MacNeil [18] and Schmidt & Phillips [35] provide tabulations for a transformed version of this statistic in the case where h_t is a linear trend. The presence of a unit root is rejected at the 5% level if $R_{VN} > cv(R_{VN}, 5\%)$.

The parametric ADF test

The most common parametric unit root test is based on the following autoregressive approximation to (4) :

$$(8) \quad \Delta y_t = \alpha y_{t-1} + \sum_{j=1}^k \varphi_j \Delta y_{t-j} + \beta' x_t + \varepsilon_t.$$

As $k \rightarrow \infty$ we can expect the autoregressive approximation to give an increasingly accurate representation of the true process. The unit root hypothesis in (4) corresponds to the hypothesis $\alpha = 0$ in (8). The hypothesis is tested by means of the regression t-ratio statistic on the coefficient α . This statistic has the same limit distribution (and critical values) as the Z_t test given in (6) above, provided $k \rightarrow \infty$ at an appropriate rate as $n \rightarrow \infty$ [32]. The test is known as the augmented Dickey-Fuller (ADF) test.

Efficient detrending by quasi-differencing

As discussed above, the von Neumann ratio/LM test R_{vN} is constructed using an efficient detrending regression under the null hypothesis in contrast to the regression (4), where there are redundant trending regressors. One way to improve the power of unit root tests is to perform the detrending regression in a way that is efficient under the alternative hypothesis as well, an idea that was suggested in [7] in the context of the removal of means and linear trends. Alternatives that are close to unity can often be well modelled using the local alternative [25]

$$(9) \quad \alpha = \exp(n^{-1}c) \sim 1 + n^{-1}c$$

for some fixed $c = \bar{c}$, say, given the sample size n . Quasi-differencing rather than differencing can now be used in the detrending regression. Such a regression leads to estimates of the trend coefficients that are asymptotically more efficient than an OLS regression in levels [16], and this result justifies the modified test procedure that follows.

Set $\Delta_{\bar{c}} y_t = (1 - L - n^{-1}\bar{c}L)y_t = \Delta y_t - n^{-1}\bar{c}y_{t-1}$, and run the detrending OLS regression $\Delta_{\bar{c}} y_t = \tilde{\gamma}' \Delta_{\bar{c}} x_t + \Delta_{\bar{c}} \tilde{y}_t^s$. Using the fitted coefficients $\tilde{\gamma}$, the levels data are detrended according to $\tilde{y}_t = y_t - \tilde{\gamma}' x_t$, and \tilde{y}_t can be used in the construction of all of the above unit root tests. For example, the modified semi-parametric Z_α test has the form $\tilde{Z}_\alpha = n(\tilde{\alpha} - 1) - \tilde{\lambda} \left(n^{-2} \sum_{t=2}^n \tilde{y}_{t-1}^2 \right)^{-1}$, where $\tilde{\lambda}$ is a consistent estimator of λ , and $\tilde{\alpha}$ is the coefficient in the regression of \tilde{y}_t on \tilde{y}_{t-1} . New critical values are needed for the \tilde{Z}_α test and the limit theory depends not only on the trend functions, as it does in (5), but also on the localizing parameter \bar{c} that is used in the quasi-

differencing. A good default choice of \bar{c} seems to be the value for which local asymptotic power is 50% [7, 14].

A point optimal test

When the model for y_t is a Gaussian AR(1) with unit error variance, the Neyman-Pearson lemma can be used to construct the most powerful test of a unit root against a simple point alternative. This is a point optimal test (POT, [14]) for a unit root at the alternative that is selected. Taking a specific local alternative with $c = \bar{c}$ in (9), using quasi-differencing to detrend, and using a consistent nonparametric estimate $\hat{\omega}^2$ of the nuisance parameter ω^2 , the POT test statistic for a unit root in (1) and (2) has the form $\tilde{P}_{\bar{c}} = \hat{\omega}^{-2} \{ \bar{c}^2 n^{-2} \sum_{t=2}^n (\tilde{y}_{t-1}^s)^2 - \bar{c} n^{-1} \tilde{y}_n^s \}$, which was given by Elliot *et al.* [7] in the case where there is a linear trend in (1). The test is performed by comparing the observed value of the statistic with the critical value obtained by simulation. The presence of a unit root in the data is rejected at the 5% level if $\tilde{P}_{\bar{c}} < cv(\tilde{P}_{\bar{c}}, 5\%)$, *i.e.* if $\tilde{P}_{\bar{c}}$ is too small. Note that in the construction of $\tilde{P}_{\bar{c}}$, the estimate $\hat{\omega}^2$ is used and this is obtained in the same way as in the Z_t test, *i.e.* using residuals from the regression (4).

Asymptotic properties and local power

The above test statistics are asymptotically similar* in the sense that their limit distributions are free of nuisance parameters. But, the limit distributions do depend on whether the data has been prefiltered in any way by a preliminary regression. The tests are also consistent against stationary alternatives provided that any nonparametric estimator of ω^2 that is used in the test converges in probability to a positive limit as $n \rightarrow \infty$. The latter condition is important, and it typically fails when estimates of ω^2 are constructed using first differences or quasi-differences of the data rather than regression residuals [27].

Rates of divergence of the statistics under the alternative are also available. For instance, when $|\alpha| < 1$, $Z_\alpha, \tilde{Z}_\alpha, R_{\nu N} = O_p(n)$, and $Z_t, ADF = O_p(n^{1/2})$ as $n \rightarrow \infty$ [27]. Thus, coefficient-based tests that rely on the estimated autoregressive coefficient and the von Neumann ratio/LM

tests diverge at a faster rate than tests that are based on the regression t-ratio. We may therefore expect such tests to have greater power than t-ratio tests, and this is generally borne out in simulations. Heuristically, the t-ratio tests suffer because there is no need to estimate a scale parameter when estimating the autoregressive coefficient α .

Under the local alternative (9), the limit theory can be used to analyse local asymptotic power. When (2) and (9) hold, y_t^s behaves asymptotically like a linear diffusion rather than Brownian motion, *i.e.* $n^{-1/2}y_{[nr]}^s \Rightarrow J_c(r) = \int_0^r e^{(r-s)c} dW(s)$. The limit distributions of the unit root test statistics then involve functionals of $J_c(r)$ [25]. The local asymptotic theory can be used to construct asymptotic power envelopes for unit root tests by taking the limit distribution of the POT statistic under the local alternative $c = \bar{c}$, and then varying the parameter \bar{c} .

Finite sample properties of unit root tests

Extensive simulations have been conducted to explore the finite sample performance of unit root tests. One general conclusion to emerge is that the discriminatory power in all of the tests between models with a root at unity and a root close to unity is low. For instance, power is less than 30% for $\alpha \in [0.90, 1.0)$ and $n = 100$. Power is reduced further by detrending the data. Both these features mirror the asymptotic theory. One interesting finding from simulation studies is the extent of the finite sample size distortion of the tests in cases where the true model is close to a trend stationary process. For example, if u_t in (2) follows a moving average process $u_t = \varepsilon_t + \theta\varepsilon_{t-1}$ with θ large and negative, then the sample trajectories of y_t^s more closely resemble those of a stationary process than a random walk. In such cases there is a tendency for all of the tests to overreject the null of a unit root. Tests that are based directly on autoregressive coefficient estimates, like Z_α , tend to be more affected by size distortion than the other tests. This is because the bias in the first order autoregressive estimator is large in this case, not only in finite samples but even in the asymptotic distribution (7), where the miscentering is measured by the bias parameter $\lambda = \theta\sigma_\varepsilon^2$. Good estimates of the bias parameter are needed to control the size

distortion. Since λ is estimated in a nonparametric way by kernel methods, it is usually not estimated at a \sqrt{n} rate. Recent attempts to improve the estimation of this parameter using data-determined bandwidth choices [1], prefiltering [2] and data-based model selection and prefiltering [16] offer some promise, the latter reference showing that \sqrt{n} rates of estimation are achievable in these estimates when consistent *model selection** techniques are used to determine the prefilter.

The parametric ADF procedure is less affected by size distortions when the true model is close to stationarity, but generally has much less power than the other tests. With this test, power is further reduced by the inclusion of additional lagged dependent regressors in (4). Again, use of model selection methods like BIC [36] are useful in this respect and these provide some increase in the finite sample power of the ADF test.

Since detrending the data reduces power, surplus trend variables in regressions like (4) will do so also. Hence, efficient detrending procedures can be expected to benefit all the tests. Simulations confirm [38] that detrending by regression in quasi-differences seems to be the most successful method so far for increasing finite sample (and asymptotic) power.

Trends with structural breaks

Breaks in deterministic trend functions are often employed to capture changes in trend. This possibility is already included in the specification of h_t in (1). For instance, the trend function

$$(10) \quad h_t = \sum_{j=0}^p f_j t^j + \sum_{j=0}^p f_{m,j} t_m^j, \quad \text{where} \quad t_m^j = \begin{cases} 0 & t \in \{1, \dots, m\} \\ (t-m)^j & t \in \{m+1, \dots, n\} \end{cases}$$

has a time polynomial of degree p (the first component) and a similar time polynomial with different coefficients (the second component) that initiates at the point $t = m+1$. This trend function therefore allows for the presence of a structural change in the polynomial trend at the data point $t = m+1$. Suppose $\mu = \lim_{n \rightarrow \infty} (m/n) > 0$ is the limit of the fraction of the sample where this structural change occurs. Then the limiting trend function $X(r)$ corresponding to (10) has a similar break at the point μ . The unit root tests given above, including those that make use of efficient detrending procedures, continue to apply for such broken trend functions. Indeed, (10)

may be extended further to allow for multiple break points in the sample and in the limit process without affecting the theory.

In order to construct unit root tests that allow for breaking trends like (10) it is necessary to specify the break point m . (Correspondingly, the limit theory depends on limit processes that depend on the break point μ). In effect, the break point is exogenously determined. Perron [23] considered linear trends with single break points in this way. An alternative perspective is that any break points are endogenous to the data and unit root tests should take account of this fact. Alternative unit root tests have been suggested [40] that endogenise the break point by choosing the value of m that gives the least favourable view of the unit root hypothesis. This has been done for the parametric ADF test and for linear trends with breaks. If $ADF(m)$ denotes the ADF statistic given by the t-ratio for α in the ADF regression (4) with a broken trend function like (10), then the trend break ADF statistic is

$$(11) \quad ADF(\hat{m}) = \min_{\bar{m} \leq m \leq \bar{\bar{m}}} ADF(m), \quad \text{where } \bar{m} = [n\bar{\mu}], \quad \bar{\bar{m}} = [n\bar{\bar{\mu}}] \quad \text{and} \quad 0 < \bar{\mu} < \bar{\bar{\mu}} < 1.$$

The limit theory for this trend break ADF statistic is given by

$$(12) \quad ADF(\hat{m}) \Rightarrow \inf_{\mu \in [\bar{\mu}, \bar{\bar{\mu}}]} \left[\int_0^1 W_X dW \right] \left[\int_0^1 W_X^2 \right]^{-1/2},$$

where the limit process $X(r)$ that appears in this functional on the right side is now dependent on the trend break point μ over which the functional is minimised. Critical values of the limiting test statistic (12) are further out in the tail than the those of the exogenous trend break statistic, so it is harder to reject the null hypothesis of a unit root when the break point is considered to be endogenous. Simulations indicate that the introduction of trend break functions leads to further reductions in the power of unit root tests. Sample trajectories of a random walk are often similar to those of a process that is stationary about a broken trend for some particular breakpoint (more so when several break points are permitted in the trend). So reductions in the power of unit root tests against competing models of this type should not be unexpected.

Seasonal unit root tests

The parametric ADF test has been extended to the case of seasonal unit roots. In order to accommodate fourth differencing the autoregressive model is written in the new form

$$(13) \quad \Delta_4 y_t = \alpha_1 y_{1t-1} + \alpha_2 y_{2t-1} + \alpha_3 y_{3t-2} + \alpha_4 y_{3t-1} + \sum_{j=1}^p \phi_j \Delta_4 y_{t-j} + \varepsilon_t$$

where $\Delta_4 = 1 - L^4$, $y_{1t} = (1 + L)(1 + L^2)y_t$, $y_{2t} = -(1 - L)(1 + L^2)y_t$, $y_{3t} = -(1 - L^2)y_t$. The data y_{1t}, y_{2t}, y_{3t} retain the unit root at the zero frequency (long run), the semi-annual frequency (two cycles per year) and the annual frequency (one cycle per year), respectively. When $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 0$, there are unit roots at the zero and all seasonal frequencies. To test the hypothesis of a unit root ($L = 1$), a t-ratio test of $\alpha_1 = 0$ is used. Similarly, the test for a semi-annual root ($L = -1$) is based on a t-ratio test of $\alpha_2 = 0$, and the test for an annual root on the t-ratios for $\alpha_3 = 0$ or $\alpha_4 = 0$. Details of the implementation of this procedure are given in Hylleberg *et al.* [12]. The limit theory is developed in Chan & Wei [3].

Bayesian tests

While most practical work on unit root testing has utilised classical procedures of the type discussed above, Bayesian methods offer certain advantages that are useful in empirical research. Foremost among these is the potential that these methods offer for embedding the unit root hypothesis in the wider context of model specification. Whether or not a model such as (4) has a unit root can be viewed as part of the bigger issue of model determination. Model comparison techniques like *posterior odds* and *predictive odds* make it easy to assess the evidence in the data in support of the hypothesis $\alpha = 1$ at the same time as decisions are made concerning other features of the model, such as the lag order in the autoregression (4), the degree of the deterministic trend component and the presence of trend breaks. Phillips & Ploberger [29, 30] explore this approach to unit root testing and give an extension of the Schwarz [36] criterion that can be used for this purpose in models with nonstationary data.

A second advantage of Bayesian methods in models with unit roots is that the asymptotic form of the posterior density is normal [13, 30] a result that facilitates large sample Bayesian inference and contrasts with the non-standard asymptotic distribution theory of classical estimators and tests. Thus, a large sample Bayesian confidence set for the autoregressive parameter α in (4) can be constructed in the conventional way without having to appeal to any nonstandard limit theory. In this respect, Bayesian theory (which leads to a symmetric confidence set for α) differs from classical statistical analysis where the construction of valid confidence regions is awkward because of the discontinuity of the limit theory at $\alpha = 1$ (but may be accomplished using local asymptotics). This divergence can lead to quite different inferences being made from the two approaches with the same data. This is so even when the influence of the prior is negligible, as it is in very large samples. In small samples, the role of the prior is important and time series models raise special concerns about the construction of uninformative priors, primarily because a great deal is known about the properties of simple time series models like autoregressions and their characteristic features in advance of data analysis. How this knowledge should be used or whether it should be ignored is a matter on which there is ongoing debate (Phillips [26] and two recent themed issues of the *Journal of Applied Econometrics*, 1991, and *Econometric Theory*, 1994).

Third, Bayesian methods offer flexibility and convenience in analysing models with possible unit roots and endogenous trend breaks. In such cases a prior distribution of break points is postulated (such as a uniform prior across potential break points), the posterior mass function is calculated, and the Bayes estimate of the break point is taken as the one with highest posterior mass [41]. This approach makes the analysis of multiple break points straightforward, a problem where classical asymptotic theory is much more complex.

Testing Stationarity

Adding a stationary component v_t to (1) and (2) gives the model

$$(14) \quad y_t = h_t + y_t^s + v_t, \quad y_t^s = y_{t-1}^s + u_t,$$

which decomposes the time series y_t into a deterministic trend, a stochastic trend and a stationary residual. The stochastic trend in (14) is annihilated when $\sigma_u^2 = 0$, which therefore corresponds to a null hypothesis of trend stationarity. Under Gaussian assumptions and iid error conditions, the hypothesis can be tested in a simple way using the LM principle, and the procedure is easily extended to more general cases where there is serial dependence, by using parametric [17], or semiparametric methods [15]. Defining $w_t = y_t^s + v_t$ and writing its differences as $\Delta w_t = (1 - \theta L)\eta_t$ where η_t is stationary, it is clear that $\sigma_u^2 = 0$ in (14) corresponds to the null hypothesis of a moving average unit root $\theta = 1$. Thus, there is a correspondence between testing for stationarity and testing for a moving average unit root [33].

Applications, empirical evidence and future prospects

Most empirical applications of unit root tests have been in the field of economics. Martingales* play a key role in the mathematical theory of efficient financial markets [6], and in the macroeconomic theory of the aggregate consumption behavior of rational economic agents [11]. In consequence, economists have been intrigued by the prospect of testing these theories. In the first modern attempt to do so using unit root tests, Nelson and Plosser [19] tested fourteen historical macroeconomic time series for the United States by the ADF test and found empirical evidence to support a unit root for thirteen of these series (the exception being unemployment). Since then, these series have been re-tested with other methods, and hundreds of other time series have been examined in the literature. While it is recognised that the discriminatory power of unit root tests is often low, there is a mounting body of evidence that many economic and financial time series are well characterised by models with roots at or near unity, as in the case of the leading economic indicators data graphed in Figure 1.

In empirical applications to multiple time series, the ADF and semi-parametric Z tests have been extensively used to test for the presence of cointegration* (or co-movement among variables with unit roots). The tests are used in the same way as unit root tests and have the same null hypothesis, but the data are the residuals from an OLS regression among the variables, and the

alternative hypothesis (of cointegration) is now the main hypothesis of interest [8, 27]. The model is analogous to (1), but both variables y_t and x_t have unit roots and y_t^s is stationary.

Unit root models, testing procedures, and unit root asymptotics now occupy a central position in the econometric analysis of time series. This is partly because of the growing empirical evidence of stochastic trends in economic data, and partly because of the importance of the notion of shock persistence in economic theory. The scope for the use of these methods in empirical research in other fields like political science and communications seems substantial. Advances in computer technology will continue to facilitate the use of simulation methods in dealing with the non standard distributions that unit root methods entail. The explosion of research over the last decade in the field of nonstationary time series and unit root methods shows no sign of abating. The field is full of potential for future developments in statistical theory, in modeling and in empirical applications.

References

- [1] Andrews, D.W.K., 1991, "Heteroskedasticity and autocorrelation consistent covariance matrix estimation", *Econometrica*, 59, 817-858.
- [2] Andrews, D.W.K. & J.C. Monahan, 1992, "An improved heteroskedasticity and autocorrelation consistent covariance matrix estimator", *Econometrica*, 60, 953-966.
- [3] Chan N.H. & C.Z.Wei, 1988, "Limiting distributions of least squares estimates of unstable autoregressive processes," *Annals of Statistics*, 16, 367-401. (Develops limit theory of least squares estimates in models with roots on the unit circle).
- [4] Dickey, D. A. & W. A. Fuller, 1979, "Distribution of estimators for autoregressive time series with a unit root", *J. Amer. Statist. Assoc.*, 74, 427-431. (Develops limit theory of tests for a Gaussian random walk.)
- [5] Dickey, D. A. & W. A. Fuller, 1981, "Likelihood ratio tests for autoregressive time series with a unit root", *Econometrica*, 49, 1057-1072.

- [6] Duffie, D., 1988, *Security Markets: Stochastic Models*. San Diego: Academic Press.
- [7] Elliot, G., T.J. Rothenberg & J.H. Stock, "Efficient tests for an autoregressive unit root", 1992, NBER working paper no.130.
- [8] Engle, R.F. & C.W.J. Granger, 1987, "Cointegration and error correction: representation, estimation and testing", *Econometrica*, 55, 251-276.
- [9] Evans, G.B.A. & N.E. Savin, 1981, "The calculation of the limiting distribution of the least squares estimator of the parameter in a random walk model", *Annals of Statistics*, 9, 1114-1118. (Graphs the limit distribution of the OLS estimate of the coefficient in a random walk.)
- [10] Grenander, U. & M. Rosenblatt, 1957, *Statistical Analysis of Stationary Time Series*, New York: John Wiley.
- [11] Hall, R.E., 1978, "Stochastic implications of the life cycle-permanent income hypothesis: theory and evidence", *J. Polit. Econom.*, 86, 971-987. (Derives a martingale theory of consumption from representative agent economic behavior).
- [12] Hylleberg, S., R.F. Engle, C.W.J. Granger & S. Yoo, 1990, "Seasonal integration and cointegration", *J. Econometrics*, 44, 215-238.
- [13] Kim, J-Y., 1994, "Bayesian asymptotic theory in a time series model with a possible nonstationary process", *Econometric Theory*, 10, 764-773.
- [14] King M.L., 1988, "Towards a theory of point optimal testing", *Econometric Reviews*, 6, 169-218.
- [15] Kwiatkowski, D., P.C.B. Phillips, P. Schmidt & Y. Shin, 1992, "Testing the null hypothesis of stationarity against the alternative of a unit root: how sure are we that economic time series have a unit root?", *J. Econometrics*, 54, 159- 178. (Derives LM test of null hypothesis of stationarity.)
- [16] Lee C.C. & P.C.B. Phillips, 1994, "Efficiency gains using GLS over OLS under nonstationarity", Yale University manuscript.

- [17] Leybourne S. J. & B.P.M. McCabe, 1994, "A consistent test for a unit root", *J. Bus. Econ. Statist.*, 12, 157-166.
- [18] MacNeill, I.B., 1978, "Properties of sequences of partial sums of polynomial regression residuals with applications to tests for change of regression at unknown times", *Annals of Statistics*, 6, 422-433.
- [19] Nelson, C.R. & C. Plosser, 1982, "Trends and random walks in macro-economic time series: some evidence and implications", *J. Monet. Econom.*, 10, 139-162. (First major empirical application of unit root tests in economics).
- [20] Ouliaris, S., J.Y. Park & P.C.B. Phillips, 1989, "Testing for a unit root in the presence of a maintained trend", pp. 7-28 in B.Raj (ed.) *Advances in Econometrics and Modelling*, Amsterdam: Kluwer.
- [21] Ouliaris, S., & P.C.B. Phillips, 1994, *Coint 2.0*, Aptech Systems, Maple Valley, WA.
(Software for unit root and cointegration testing).
- [22] Park, J.Y. & J. Sung, 1994, "Testing for unit roots in models with structural change", *Econometric Theory*, 10, 917-936.
- [23] Perron, P., 1989, "The great crash, the oil price shock and the unit root hypothesis", *Econometrica*, 57, 1361-1401. (Derives tests for a unit root against stationarity about a breaking trend).
- [24] Phillips, P.C.B., 1987, "Time series regression with a unit root", *Econometrica*, 55, 277-302. (Develops semi-parametric versions of the Dickey-Fuller unit root tests and shows how to derive asymptotics using functional central limit theory).
- [25] Phillips, P.C.B., 1987, "Towards a unified asymptotic theory of autoregression", *Biometrika*, 74, 535-547. (Develops local power asymptotics for unit root tests).
- [26] Phillips, P.C.B., 1991, "To criticize the critics: an objective Bayesian analysis of stochastic trends", *J. Applied Econometrics*, 6, 333-364. (Studies uninformative priors for possibly nonstationary autoregressions).

- [27] Phillips, P.C.B. & S. Ouliaris, 1990, "Asymptotic properties of residual based tests for cointegration", *Econometrica*, 58, 165-194. (Applies unit root tests to test for cointegration and derives limit distribution theory of tests.)
- [28] Phillips, P.C.B. & P. Perron, 1988, "Testing for unit roots in time series regression", *Biometrika*, 75, 335-346.
- [29] Phillips, P.C.B. & W. Ploberger, 1994, "Posterior odds testing for a unit root with data-based model selection", *Econometric Theory*, 10, 774-808. (Bayesian model selection principle to test for a unit root in conjunction with ARMA lag orders and trend degree.)
- [30] Phillips, P.C.B. & W. Ploberger, 1995, "An asymptotic theory of Bayesian inference for time series", *Econometrica*, in press.
- [31] Phillips, P. C. B. & V. Solo, 1992, "Asymptotics for linear processes", *Annals of Statistics*, 20, 971-1001.
- [32] Said, S.E. & D.A. Dickey, 1984, "Testing for unit roots in autoregressive-moving average models of unknown order", *Biometrika*, 71, 599-608. (Derives asymptotic theory for the augmented Dickey-Fuller test).
- [33] Saikkonen, P. & R. Luukkonen, 1993, "Testing for a moving average unit root in autoregressive integrated moving average models", *J. Amer. Statist. Assoc.*, 88, 596-601.
- [34] Sargan, J.D. & A. Bhargava, 1983, "Testing residuals from least squares regression for being generated by the Gaussian random walk", *Econometrica*, 51, 153-174. (Suggests von Neumann ratio tests for a Gaussian random walk.)
- [35] Schmidt, P. & P.C.B. Phillips, 1992, "Testing for a unit root in the presence of deterministic trends", *Oxford Bulletin of Economics and Statistics*, 54, 257-288. (Derives LM test of a unit root in models with deterministic trends.)
- [36] Schwarz, G., 1978, "Estimating the dimension of a model", *Annals of Statistics*, 6, 461-464.
- [37] Solo, V., 1984, "The order of differencing in ARIMA models", *J. Amer. Statist. Assoc.*, 79, 916-921.

- [38] Stock, J.H., 1995, "Unit roots, structural breaks and trends", in R.F. Engle & D. McFadden (eds.), *Handbook of Econometrics Vol. 4*, Amsterdam: North Holland. (Review article of research on unit roots, trends and structural change up to the early 1990's.)
- [39] White, J. S., 1958, "The limiting distribution of the serial correlation coefficient in the explosive case", *Annals of Math. Statist.*, 29, 1188-1197.
- [40] Zivot, E. & D.W.K. Andrews, 1992, "Further evidence on the great crash, the oil price shock, and the unit root hypothesis", *J. Bus. Econ. Statist.*, 10, 251-270. (Develops limit theory of unit root tests against models with unknown structural breaks).
- [41] Zivot, E. & P.C.B. Phillips, 1994, "A Bayesian analysis of trend determination in economic time series", *Econometric Reviews*, 13, 291-336. (Bayesian analysis of unit roots, trends and trend breaks.)

Bibliography (books and additional review articles)

- Banerjee, A., J. Dolado, J. W. Galbraith & D. F. Hendry, 1992, *Cointegration, Error Correction, and the Econometric Analysis of Non-Stationary Time Series*, Oxford: Oxford University Press. (Introduction to unit roots and cointegration, designed to be accessible to graduate students of economics.)
- Campbell, J.Y. & P. Perron, 1991, "Pitfalls and opportunities: what macroeconomists should know about unit roots" *NBER Macroeconomics Annual*, 141-200. (Survey of unit roots and cointegration intended for applied researchers.)
- Diebold, F.X. & M. Nerlove, 1990, Unit roots in economic time series, *Advances in Econometrics*, 8, 3-70. (Survey of unit root literature in the 1980's.)
- Dolado, J., T. Jenkinson & S. Sosvilla-Rivero, 1990, Cointegration and unit roots. *J. Economic Surveys*, 4, 249-273. (General survey of literature in the field in the 1980's.)
- Econometric Theory*, 10, No. 3&4, 1994 (Symposium of recent work in the field of Bayesian unit roots.)

Econometric Reviews 1994 (Two special issues on unit roots and cointegration reporting recent research.)

Fuller, W.A. , 1976, *Introduction to Statistical Time Series*, New York: Wiley. (Highly accessible technical introduction to time series. The first textbook treatment of unit roots. Contains the original tabulations of the critical values of the Dickey-Fuller tests.)

Hamilton, J.D., 1994, *Time Series Analysis*, Princeton: Princeton University Press. (Recent textbook treatment of unit root theory and cointegration with examples and applications. Designed mainly for graduate students in economics.)

Journal of Applied Econometrics, 6, No. 4, 1991. (Special issue on Bayesian unit root models with focus on the formulation of priors in stationary and nonstationary time series models.)

Journal of Business and Economic Statistics, 10, 1992. (Special issue devoted to structural change models and unit roots with broken trends.)

Oxford Bulletin of Economics and Statistics, vols. 48 & 54, 1986 & 1992 (Two special issues of empirical and theoretical articles on cointegration and unit roots.)

Phillips, P.C.B., 1988, Multiple regression with integrated time series (Detailed review article on the asymptotic theory of unit roots, regression with integrated processes, spurious regression, cointegration, and local-to-unity asymptotics.)

Phillips, P.C.B., 1992, Unit roots, *New Palgrave Dictionary of Money and Finance*, London: Macmillan. (Short introduction to unit root testing and unit root models in economics and finance.)

Phillips, P.C.B., 1995, Cointegration and unit roots: recent books and themes for the future.

Journal of Applied Econometrics, 10, 87-94. (Reviews recent themes in the literature and points to some directions of future research.)

Cross References

Autocorrelation; Brownian motion; Brownian bridge; cointegration; differencing; kernel estimation; martingale; model choice; nonstationarity; random walk; seasonal model; similar test; spectral density; stationarity; trend;

List of Unit Root Tests

Augmented Dickey-Fuller (ADF) test; Dickey-Fuller test; LM test; Phillips Z-tests; Point optimal test; von Neumann ratio test.

Figure 1: Monthly U.S. Economic Time Series
1948:1 - 1994:1




Figure 2: Unit Root Limit Densities

